arXiv:1505.07597v1 [cond-mat.soft] 28 May 2015

One-dimensional Kac model of dense amorphous hard spheres

H. Ikeda[1] and A. Ikeda[2]

[1] *Department of Physics, Nagoya University - Nagoya, 464-8602, Japan*
[2] *Fukui Institute for Fundamental Chemistry, Kyoto University - Takano-Nishihiraki-cho 34-4, Sakyo-ku, Kyoto, 606-8103, Japan*

PACS `64.70.Q` – Theory and modeling of the glass transition
PACS `05.20.-y` – Classical statistical mechanics.
PACS `83.80.Fg` – Granular solids.

Abstract – We introduce a new model of hard spheres under confinement for the study of the glass and jamming transitions. The model is an one-dimensional chain of the d-dimensional boxes each of which contains the same number of hard spheres, and the particles in the boxes of the ends of the chain are quenched at their equilibrium positions. We focus on the infinite dimensional limit ($d \to \infty$) of the model and analytically compute the glass transition densities using the replica liquid theory. From the chain length dependence of the transition densities, we extract the characteristic length scales at the glass transition. The divergence of the lengths are characterized by the two exponents, $-1/4$ for the dynamical transition and -1 for the ideal glass transition, which are consistent with those of the p-spin mean-field spin glass model. We also show that the model is useful for the study of the growing length scale at the jamming transition.

Introduction. – When liquids are rapidly cooled well below the melting temperature, the relaxation time and viscosity drastically increase and eventually they freeze into the disordered state. This phenomenon is known as the glass transition [1]. The mechanism of this slowing down and the presence/absence of an underlying genuine phase transition are under active debate [2–4].

A good way to understand theoretically the glass transition would be to first construct a mean field theory and then consider finite dimensional effects. Recently the glass transition of the infinite dimensional hard spheres is studied as a mean-field model, and the exact thermodynamic theory of the model is developing [5–8]. The glass transition of the model can be characterized by the two relevant densities [5]. The first is the dynamical transition density, φ_d, below which the phase space is divided into an exponentially large number of metastable states. In the mean-field limit, the dynamics of the system gets frozen at this density as the system is trapped into one of the metastable states. With increasing the density, the number of the metastable states decreases and eventually becomes sub-exponential, at which the system undergoes the thermodynamic transition into the ideal glass state. This transition density φ_K is called the Kauzman density.

What types of finite dimensional effects play a dominant role in the glass transition is not clear yet [3, 9]. One promising theoretical scenario which incorporates an effect of fluctuations of finite dimensional systems is proposed as the Random First Order Transition (RFOT) theory [10]. This theory assumes that the divergence of the relaxation time at φ_d is avoided, and at φ between φ_d and φ_K, the liquid state can be seen as a patchwork of the local metastable configurations [11]. The theory predicts that the characteristic size of these domains increases with increasing the density and eventually diverges at φ_K, where the thermodynamic glass transition occurs.

In practice, this length can be measured through the thermodynamic behavior of liquids under confinement [11]. We prepare an equilibrium configuration of particles and pin only the particles outside of a cavity of the size L. One can define and compute the critical cavity size at which the boundary affects the thermodynamics of the system in the cavity, which is referred to as the point-to-set (PS) correlation length. Numerical simulations of realistic liquid models under this confinement have been carried out [12–14], which confirmed that the PS length indeed grows as the systems slow down. However, computing theoretically the PS length of realistic liquids is still very challenging [15, 16]. One of the difficulties is due to the inhomogeneous nature of the confined liquids, which requires the inhomogeneous version of the liquid state theory [17]. The p-spin mean-field spin glass model under confinement has been extensively studied [18, 19]. The studies showed that the PS length of the model diverges at the Kauzuman transition point with the power-law behavior with the exponent -1 and another length characterizing the local stability of the metastable states diverges at the dynamical transition point with the exponent $-1/4$ [18].

The main purpose of the present letter is to propose a new model of hard spheres under confinement, which can be analyzed using the *homogeneous* liquid state theory [20]. Focusing on the large dimensional limit of the model, we compute analytically the PS length using the replica liquid theory.

Since the model proposed is not a spin model but hard spheres, the model undergoes another type of the phase transition, called the jamming transition, at the higher density [5, 21]. This is the transition of the liquid state into the state where hard spheres are packed so closely that the pressure diverges. We additionally show that the model can be used to study the growing lengths at the jamming transition.

Model. – Consider an one-dimensional chain of $L+2$ boxes. Each box is a d-dimensional cube of the volume V and contains N hard spheres. We assume that a particle interacts with the other particles in the same box and those in the nearest two boxes. Therefore, the Hamiltonian of the model is

$$H = \sum_{l=0}^{L+1}\sum_{i<j}^{N} v(|x_i^l - x_j^l|) + \sum_{l=0}^{L}\sum_{ij}^{N} v(|x_i^l - x_j^{l+1}|), \qquad (1)$$

where the d-dimensional vector x_i^l denotes the position of the i-th particle in the l-th box and $v(r)$ is the interaction potential between hard spheres of the diameter D, hence $v(r) = \infty \ (r \le D),\ 0 \ (r > D)$. We study the model under the condition that the particles in the 0-th and $(L+1)$-th boxes are quenched at their equilibrium positions. Thus the model is interpreted as the d-dimensional hard spheres confined by the amorphous walls. We sketch a typical configuration of our model schematically in Fig. 1.



Fig. 1: A typical configuration of the model: The position of the particles in each boxes labeled 0-th to $L+1$-th are schematically shown as cycles. The shaded cycles represent the frozen particles in 0-th and $(L+1)$-th boxes, and the filled cycles represent the mobile particles in each box. In the l-th box, the particle positions in the $(l-1)$-th and $(l+1)$-th box are also shown as the solid and dashed cycles respectively to indicate the particles in l-th box can not overlap with the particles in $(l-1)$-th and $(l+1)$-th boxes.

As it is clear from the Hamiltonian, eq. (1), we can treat the model as a $L+2$ component *homogeneous* liquid, which enables us to analyze the model using the *homogeneous* liquid state theory. This is one of the advantages of this model. In particular, the free energy of this model can be evaluated analytically at the large dimensional limit using the replica liquid theory. In the following sections, we calculate the two relevant densities φ_d and φ_K for the glass transition as a function of the number of the boxes L. By interpreting L as the correlation length, we extract the lengths ξ_d and ξ_K which diverge at φ_d and φ_K, respectively. Additionally, we also apply this method to the jamming transition to discuss the possibility to compute its growing length scales.

Free energy at large dimensional limit. – To compute the free energy of the model, we employ the replica liquid theory assuming the one-step replica symmetry breaking (1RSB) ansatz [5, 22, 23]. The main idea of the theory is to consider m copies (replicas) of the original system. There is no interaction between the replicas, and therefore the partition function of the replicated system is written as

$$Z_m \equiv \prod_{a=1}^{m} \mathrm{Tr}_{x_a} e^{-\sum_{a=1}^{m} H[\{x_a\}]/T}, \qquad (2)$$

where H is the Hamiltonian of the original system, and T is the temperature. From now on, we set $T=1$ because the temperature is an irrelevant variable for hard spheres. Within the replica theory, the logarithm of the number of the minima, which is called the configurational entropy, Σ, can be calculated by the formula [23]:

$$\Sigma = -\frac{m^2}{N}\frac{\partial}{\partial m}\left(\frac{\log Z_m}{m}\right). \qquad (3)$$

If one takes the large dimensional limit ($d \to \infty$) and the thermodynamic limit ($N \to \infty$, $V \to \infty$, N/V=const), the calculation of the partition function of the replicated system can be greatly simplified because the higher order terms of the Mayer cluster expansion become negligible, and only the first term needs to be considered [5, 20]. The replicated free-energy of our model is

$$\begin{aligned}
\log Z_m = &\sum_{l=1}^{L}\int d\overline{x}\rho_l(\overline{x})\left(1-\log\rho_l(\overline{x})\right) \\
&+\frac{1}{2}\sum_{l=1}^{L}\int d\overline{x}d\overline{y}\rho_l(\overline{x})\rho_l(\overline{y})f(\overline{x}-\overline{y}) \\
&+\sum_{l=1}^{L-1}\int d\overline{x}d\overline{y}\rho_l(\overline{x})\rho_{l+1}(\overline{y})f(\overline{x}-\overline{y}) \\
&+\int d\overline{x}dy\rho_1(\overline{x})\hat{\rho}_0(y)f(\overline{x}-y) \\
&+\int d\overline{x}dy\rho_L(\overline{x})\hat{\rho}_{L+1}(y)f(\overline{x}-y), \qquad (4)
\end{aligned}$$

where $\overline{x} = \{x^1, \cdots x^m\}$ denotes the positions of hard spheres in the replicated space, and $\rho_l(\overline{x}) = \left\langle \sum_i \prod_{a=1}^m \delta(x^a - x_i^{l,a}) \right\rangle$ is the density distribution of the l-th box. The boundary conditions are included in the last two lines of eq. (4), where $\hat{\rho}_0(x) = \sum_i \delta(x - x_i^0)$ and $\hat{\rho}_{L+1}(x) = \sum_i \delta(x - x_i^{L+1})$ are the microscopic density distributions of the frozen particles belonging in the 0-th and $(L+1)$-th boxes. x_i^0 and x_i^{L+1} are taken from their equilibrium positions. If the system is sufficiently large, the self-averaging properties for the free-energy holds:

$$\log Z_m \approx \overline{\log Z_m}, \tag{5}$$

where the overline denotes the average of $\hat{\rho}_0(x)$ and $\hat{\rho}_{L+1}(x)$. Since the free-energy eq. (4) depends linearly on those variables, we have only to replace as $\hat{\rho}_0(x) \to \rho$ and $\hat{\rho}_{L+1}(x) \to \rho$, where $\rho = N/V$ is the number density of each boxes.

In the next step, we introduce the 1RSB Gaussian ansatz [5]:

$$\rho_l(\overline{x}) = \rho \int dX \prod_{a=1}^m \gamma_{A_l}(x_a - X), \tag{6}$$

where $\gamma_A(x) = \exp\left(-x^2/2A\right)/(2\pi A)^{d/2}$. This ansatz claims that the distribution of the replicated particles is the Gaussian with the variance $A_l = \left\langle \sum_{a<b}(x^{l,a} - x^{l,b}\right\rangle / m(m-1)$. Substituting eq. (6) into eq. (4) and taking the large dimensional limit [5], one obtains the analytical expression of the free energy:

$$\begin{aligned} \frac{\log Z_m}{N} &= S_{id} + S_{int}, \\ S_{id} &= \sum_{l=1}^{L} \frac{d}{2}\left[(m-1)\log\frac{\hat{A}_l}{d^2} + \log\frac{m}{d} + m\right], \\ S_{int} &= -\frac{3L\hat{\varphi}}{2}\left(1 + \frac{2}{3L}\right) + \frac{\hat{\varphi}}{2}\sum_{l=1}^{L} G_m(\hat{A}_l) \\ &+ \hat{\varphi}\sum_{l=0}^{L} G_m\left(\frac{\hat{A}_l + \hat{A}_{l+1}}{2}\right), \end{aligned} \tag{7}$$

where we introduced the normalized volume fraction $\hat{\varphi} = 2^d\varphi/d$ and the normalized cage size $\hat{A}_l = A_l d^2/D^2$. $G_m(\hat{A})$ is the auxiliary function given by

$$G_m(\hat{A}) = \int_{-\infty}^{\infty} dy e^y \left[\Theta\left(\frac{y + \hat{A}}{\sqrt{4\hat{A}}}\right)^m - \theta(y)\right], \tag{8}$$

where $\Theta(x) = [\mathrm{erf}(x) + 1]/2$. In eq. (7), the cage size of the 0-th and $(L+1)$-th boxes are zero, $\hat{A}_0 = \hat{A}_{L+1} = 0$, because of the boundary condition. $\hat{A}_l$ for other l is calculated from the saddle point conditions for $\hat{A}_l$:

$$\frac{1}{\hat{A}_l} = \frac{2}{(1-m)d}\frac{\partial S_{int}}{\partial \hat{A}_l}, \qquad (l = 1, \ldots, L). \tag{9}$$

Substituting the solution of this set of equations into eq. (7), the free energy of the m replicated system is obtained.



Fig. 2: The density dependence of the cage size for several L's. The solid line is for $L = 2$, the dashed line is for $L = 4$ and the dotted line is for $L = 8$.

Glass transition. – Here, we evaluate the correlation length of the glass transition. As mentioned before, there are the two relevant densities for the glass transition, $\hat{\varphi}_d$ and $\hat{\varphi}_K$. Their values can be computed by analyzing the free energy at $m = 1$.

First, we examine the behavior of our model near the dynamical transition point, $\hat{\varphi}_d$, where the exponentially many metastable states emerge on the free energy. In the framework of the replica liquid theory, the order parameter to characterize the metastable states is the cage size, which is given by the solutions of eq. (9) [5]. Their numerical solution for $\hat{A}_l$ of the $L/2$-th box, which is the farthest from the boundaries, are shown in Fig. 2. For the small densities, the cage size $\hat{A}_{L/2}$ is infinity $(1/(1+\hat{A}_{L/2}) = 0)$, meaning that the system is ergodic and in the liquid phase. Increasing the density, the cage size jumps discontinuously to a finite value at the dynamic transition point, $\hat{\varphi}_d$.

From Fig. 2, it is clear that $\hat{\varphi}_d$ increases as increasing L. From this result, we can convert $\hat{\varphi}_d(L)$ to $L(\hat{\varphi})$, the characteristic number of the boxes as a function of the transition density. We define the correlation length by $\xi_d = L(\hat{\varphi})/2$ and plot it in Fig. 3 [18]. We find that ξ_d diverges at $\hat{\varphi}_d$ as $\xi_d \approx (\hat{\varphi}_d - \hat{\varphi})^{-1/4}$.

Next, we evaluate the correlation length near the Kauzman density by calculating the configurational entropy, Σ, which is obtained by plugging the free energy into eq. (3). The final expression in the large dimensional limit becomes

$$\Sigma(m, \hat{\varphi}, L) = \left[\frac{d}{2}\log d - \frac{3\hat{\varphi}}{2}\left(1 + \frac{2}{3L}\right)\right] + O(d), \tag{10}$$

up to the order of $O(d \log d)$. From this expression, it is clear that Σ vanishes at

$$\hat{\varphi}_K(L) = \frac{d\log d}{3(1 + 2/3L)}. \tag{11}$$

Following the same argument as for ξ_d, the growing length



Fig. 3: The correlation legnths near the $\hat{\varphi}_d$ and $\hat{\varphi}_{th}$. The filled cycles are for ξ_d as a function of $\varepsilon = (\hat{\varphi}_d - \hat{\varphi})/\hat{\varphi}_d$. The filled squares are for ξ_{th} as a function of $\varepsilon = (\hat{\varphi}_{th} - \hat{\varphi})/\hat{\varphi}_{th}$.

around the Kauzman density is

$$\xi_K = \xi_{GCP} = \frac{L}{2} = \frac{\hat{\varphi}}{3(\hat{\varphi}_K - \hat{\varphi})}, \tag{12}$$

where $\hat{\varphi}_K = d \log d/3$ is the Kauzman density of the bulk system ($L \to \infty$).

Jammintg transition. – Hard spheres also undergo the jamming transition when the system is compressed quickly or making the pressure infinity [21]. The infinite dimensional hard spheres serve as a mean-field model of the jamming transition as well and are studied extensively using the replica liquid theory. The theory showed that the jamming transition is also characterized by the two relevant densities [5]. The jamming state obtained by compressing the ideal glass state up to the infinite pressure is referred to as the glass close packing, and its density is denoted as $\hat{\varphi}_{GCP}$. If the jammed stated is prepared by a fast compression of low density hard spheres, the pressure becomes infinity at a much lower density than $\hat{\varphi}_{GCP}$ [24]. The lowest density of the jammed states $\hat{\varphi}_{th}$ is called the threshold density.

In this section, we show that the growing length scales at the jamming transition can be computed from the analysis of the jamming transition of the present model. We essentially follow the strategy developed in the previous section. Using the replica liquid theory with the 1RSB ansatz, we compute φ_{th} and φ_{GCP} of the model as a function of the number of the boxes, L, which naturally gives rise to the two characteristic lengths $\xi_{th}(\hat{\varphi})$ and $\xi_{GCP}(\hat{\varphi})$. Note that recent studies showed that the full RSB ansatz is needed for the fully exact computation near the jamming transition [7, 8]. However such a computation for the present model seems quite involved. Here, we wish to stick to the 1RSB ansatz and demonstrate that the replica theory analysis of the model provides the analytical expressions of the growing length scales at the jamming transition.

In the replica liquid theory, the divergence of the pressure corresponds to take the $m \to 0$ limit, because m is inversely proportional to the pressure [5]. Thus $\hat{\varphi}_{th}$ can be calculated by taking the $m \to 0$ limit in the self-consistent equation for the order parameter eq. (9). Since the cage size, $\hat{A}_l$, vanishes at the $m \to 0$ limit, we set $\hat{A}_l = m\alpha_l$ [5]. This is substituted to eq. (9) before the $m \to 0$ limit is taken. We numerically solve the equation and find that the behavior of $\alpha_{L/2}$ is qualitatively the same as that of $A_{L/2}$ for the glass transition (Fig. 2). At low densities, $\alpha_{L/2} = \infty$ is the only solution of eq. (9). This means that there are no jammed states at those densities. At higher densities, the solution of eq. (9) becomes finite. This transition density is the lowest density, $\hat{\varphi}_{th}$, of the jammed states. As in the case of the glass transition, the characteristic length $\xi_{th}(\hat{\varphi})$ is computed from the L dependence of $\hat{\varphi}_{th}$. This length ξ_{th} is also plotted in Fig. 3. We find the power-law divergence $\xi_{th} \approx (\hat{\varphi}_{th} - \hat{\varphi})^{-1/4}$. Likewise, we analyze the correlation length near $\hat{\varphi}_{GCP}$. The logarithm of the number of the jammed states is calculated by the $m \to 0$ limit of the configurational entropy eq. (10) and $\hat{\varphi}_{GCP}$ is defined as the density where the configurational entropy becomes zero. From eq. (10), it is clear that the behavior of $\hat{\varphi}_{GCP}$ should be the same as those of $\hat{\varphi}_K$ because eq. (10) is independent from m. Thus we obtain the critical behavior of the diverging length near $\hat{\varphi}_{GCP}$ as $\xi_{GCP} \approx (\hat{\varphi}_{GCP} - \hat{\varphi})^{-1}$.

Summary and discussion. – In this letter, we considered an one-dimensional chain of the d-dimensional boxes each of which contains N hard spheres as a model system of liquids under confinement. By focusing on the large dimensional limit, we analytically computed the phase diagram of the model. From the chain length dependence of the transition densities, we derived the critical behavior of the two relevant length scales of the glass transition. We also showed that there are the two relevant length scales of the jamming transition, whose critical behaviors are similar to those of the glass transition as long as the 1RSB ansatz is assumed. One of the advantage of the model is that, although the system is under confinement, it can be regarded as a homogeneous liquid, and thus the model can be analyzed using the usual replica liquid theory.

We summarize the results in Fig. 4. The upper panel is the phase diagram of the glass transition of the confined liquid, where $\hat{\varphi}$ is the normalized density $\hat{\varphi} = 2^d\varphi/d$, and L is the distance between the boundaries. There are the two relevant length scales ξ_d and ξ_K, which diverge as $\xi_d \approx (\hat{\varphi}_d - \hat{\varphi})^{-1/4}$ and $\xi_K \approx (\hat{\varphi}_K - \hat{\varphi})^{-1}$, respectively. When $L > \xi_d$, the system holds the ergodicity and is in the liquid phase. For $\xi_K < L < \xi_d$, the phase space splits into the many sub-spaces and the ergodicity of the system is broken at least in the large dimensional limit. In $L < \xi_K$ region, the number of the sub-spaces becomes sub-exponential and no longer contributes to the entropy. Thus the equilibrium phase transition from the


$\xi_d \approx (\hat{\varphi}_d - \hat{\varphi})^{-1/4}$
$\xi_K \approx (\hat{\varphi}_K - \hat{\varphi})^{-1}$
L
$\hat{\varphi}$
$\xi_{th} \approx (\hat{\varphi}_{th} - \hat{\varphi})^{-1/4}$
$\xi_{GCP} \approx (\hat{\varphi}_{GCP} - \hat{\varphi})^{-1}$
No solutions
L
No solutions
$\hat{\varphi}$

Fig. 4: The schematic phase diagrams of our model: The upper and lower panel is the phase diagram for the glass and jamming transition, respectively. $\hat{\varphi} = 2^d \varphi/d$ is the normalized volume fraction and L is the confinement length. The insets are schematic pictures of the phase space in each phase, where the cycles represent the accessible regions in the phase space.

liquid phase to the ideal glass phase occurs. These results are consistent with the results for the p-spin mean-field spin glass model [18,19], where the critical behaviors $\xi_d \approx (T - T_d)^{-1/4}$ and $\xi_K \approx (T - T_K)^{-1}$ are observed. This confirms that the proposed model is considered to be a hard sphere extension of the p-spin mean-field spin glass model under confinement. The result for ξ_d is also consistent with the prediction of the inhomogeneous mode-coupling theory for the dynamic correlation length diverging at the mode-coupling transition point [17].

The lower panel of Fig. 4 is the phase diagram of the jamming transition. There are the two relevant length scales ξ_{th} and ξ_{GCP} which diverge as $\xi_{th} \approx (\hat{\varphi}_{th} - \hat{\varphi})^{-1/4}$ and $\xi_{GCP} \approx (\hat{\varphi}_{GCP} - \hat{\varphi})^{-1}$, respectively. When $L > \xi_{th}$, there are no jammed states. For $\xi_{GCP} < L < \xi_{th}$, there exist exponentially many jammed states. The number of the jammed states decreases with L, and becomes subexponential when $L < \xi_{GCP}$.

We should emphasize that the computation of ξ_{th} and ξ_{GCP} in this work is not exact, since the 1RSB ansatz is recently found to give unstable solution near the $m \to 0$ limit [7]. If we use a better ansatz, the critical exponents may change [8]. Regarding this point, we wish to indicate the two points: (1) Even though the solution is unstable, the physics of the 1RSB solution can still survive in certain situations [25] and thus the analysis given here is still useful [1]. (2) In case of ξ_{GCP}, the results would not be effected by this instability at least in the large dimensional limit, because the expression of the configurational entropy is m independent.

Finally we discuss the physical meaning of ξ_{th} and ξ_{GCP}. Several different types of lengths are known to diverge at the jamming transition and the relations between the lengths are still not clear enough [26–36]. Since ξ_{th} and ξ_{GCP} represent the distance between boundaries at which the jammed configurations first appear/disappear, we expect that these lengths correspond to the length detected in the finite size scaling analysis $\xi_{FS} \sim |\varphi - \varphi_J|^{-1.09}$ [32]. However this is a tentative expectation, because we applied a specific boundary condition (amorphous boundary) which is suitable to compute the PS length at the glass transition. It is not clear which length should be detected by this condition for the jamming transition. To resolve this question, it should be useful to study the present model under various different types of confinements such as the flat walls [37] or random boundary [19].

* * *

We thank K. Miyazaki, H. Yoshino, G. Biroli and P. Urbani for helpful discussions. HI acknowledge the JSPS Core-to-Core program and Program for Leading Graduate Schools "Integrative Graduate Education and Research in Green Natural Sciences", MEXT, Japan. AI acknowledges JSPS KAKENHI No. 26887021.

REFERENCES

[1] Debenedetti P. G. and Stillinger F. H., *Nature*, **410** (2001) 259.

[2] Götze W., *Complex Dynamics of Glass-Forming Liquids: A Mode-Coupling Theory*, Vol. **143** (Oxford University Press, Oxford) 2008

[3] Berthier L. and Biroli G., *Rev. Mod. Phys.*, **83** (2011) 587.

[4] Tanaka H., Kawasaki T., Shintani H. and Watanabe K., *Nat. Mater.*, **9** (2010) 324.

[5] Parisi G. and Zamponi F., *Rev. Mod. Phys.*, **82** (2010) 789.

[6] Kurchan J., Parisi G. and Zamponi F., *J. Stat. Mech: Theory Exp.*, **2012** (2012) P10012.

[7] Kurchan J., Parisi G., Urbani P. and Zamponi F., *J. Phys. Chem. B*, **117** (2013) 12979.

[8] Charbonneau P., Kurchan J., Parisi G., Urbani P. and Zamponi F., *J. Stat. Mech: Theory Exp.*, (2014) P10009.

[9] Biroli G. and Garrahan J. P., *J. Chem. Phys.*, **138** (2013) 12A301.

[10] Kirkpatrick T., Thirumalai D. and Wolynes P. G., *Physical Review A*, **40** (1989) 1045.

[11] Bouchaud J.-P. and Biroli G., *J. Chem. Phys.*, **121** (2004) 7347.

---

[1] For examples, if the system is not quenched well into a single inherent state during the compression and explores several different inherent states, the physics of the 1RSB solution may appear [25].

[12] Biroli G., Bouchaud J.-P., Cavagna A., Grigera T. and Verrocchio P., *Nat. Phys.*, **4** (2008) 771.

[13] Kob W., Roldán-Vargas S. and Berthier L., *Nat. Phys.*, **8** (2012) 164.

[14] Hocky F. M., Markland T. E. and Reichman D. R., *Phys. Rev. Lett.*, **108** (2012) 225506.

[15] Franz S., Jacquin H., Parisi G., Urbani P. and Zamponi F., *J. Chem. Phys.*, **138** (2013) 12A540.

[16] Cammarota C. and Seoane B., *First principle computation of random pinning glass transition, glass cooperative length-scales and numerical comparisons* arXiv:1403.7180.

[17] Biroli G., Bouchaud J.-P., Miyazaki K. and Reichman D. R., *Phys. Rev. Lett.*, **97** (2006) 195701.

[18] Franz S. and Montanari A., *J. Phys. A: Math. Theor.*, **40** (2007) F251.

[19] Cammarota C., Gradenigo G. and Biroli G., *Phys. Rev. Lett.*, **111** (2013) 107801.

[20] Hansen J.-P. and McDonald I. R., *Theory of simple liquids* (Access Online via Elsevier) 1990.

[21] Torquato S., Truskett T. M. and Debenedetti P. G., *Phys. Rev. Lett.*, **84** (2000) 2064.

[22] Mézard M. and Parisi G., *Phys. Rev. Lett.*, **82** (1999) 747.

[23] Monasson R., *Phys. Rev. Lett.*, **75** (1995) 2847.

[24] Chaudhuri P., Berthier L. and Sastry S., *Phys. Rev. Lett.*, **104** (2010) 165701.

[25] Yoshino H. and Zamponi F., *Phys. Rev. E*, **90** (2014) 022302.

[26] O'Hern C. S., Silbert L. E., Liu A. J. and Nagel S. R., *Phys. Rev. E*, **68** (2003) 011306.

[27] Silbert L. E., Liu A. J. and Nagel S. R., *Phys. Rev. Lett.*, **95** (2005) 098301.

[28] Wyart M., Silbert L., Nagel S. and Witten T., *Phys. Rev. E*, **72** (2005) 051306.

[29] Ellenbroek W. G., Somfai E., van Hecke M. and van Saarloos W., *Phys. Rev. Lett.*, **97** (2006) 258001.

[30] Wyart M., *EPL*, **89** (2010) 64001.

[31] van Hecke M., *J. Phys.: Cond. Mat.*, **22** (2010) 033101.

[32] Vågberg D., Valdez-Balderas D., Moore M. A., Olsson P. and Teitel S., *Phys. Rev. E*, **83** (2011) 030303.

[33] Ikeda A., Berthier L. and Biroli G., *J. Chem. Phys.*, **138** (2013) 12A507.

[34] Goodrich C. P., Ellenbroek, W. G. and Liu A. J., *Soft Matter*, **9** (2013) 10993.

[35] Lerner E., DeGiuli E. During G. and Wyart M., *Soft Matter*, **10** (2014) 5085.

[36] DeGiuli E., Lerner E., Brito C. and Wyart M., *Proc. Natl. Acad. Sci. USA*, **111** (2014) 17054.

[37] Desmond K. W. and Weeks E. R., *Phys. Rev. E*, **80** (2009) 051305.